

**Application
for
United States Letters Patent**

0000000000000000000000000000000000

To all whom it may concern:

**Be it known that (we) John F. O'Connor, Galina I. Kovalevskaya, and
Steven Birken**

have invented certain new and useful improvements in

METHODS FOR PREDICTING PREGNANCY OUTCOME IN A SUBJECT BY hCG ASSAY

of which the following is a full, clear and exact description.

METHODS FOR PREDICTING PREGNANCY OUTCOME
IN A SUBJECT BY hCG ASSAY

Handwritten initials "AS AI" inside a large handwritten checkmark.

This application is a continuation-in-part application of U.S. Serial No. 09/017,976, filed February 3, 1998, the contents of which are hereby incorporated by reference into this application.

5

The invention disclosed herein was made with United States Government support under National Institutes of Health Grant Nos. NIEHS ES-07589 and HD 15454. Accordingly, the U.S. Government has certain rights in this invention.

10

Background of the Invention

Throughout this application, various publications are referenced by author and date. Full citations for these publications may be found listed alphabetically at the end of the specification immediately preceding the claims. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art.

15

20

Early pregnancy loss (EPL) is a widespread, but largely undiagnosed problem. In order to adequately diagnose and develop treatments for EPL it is essential to be able to detect and measure the rate of occurrence of EPL. This is critically important in epidemiological studies, some of which are related to exposures to known or suspected reproductive toxins in the workplace, in the environment or by personal use. These early pregnancy losses are often not recognized by women or physicians and are detected solely by the measurement of hCG in the urine at the time between implantation and expected menses. They are sometimes termed "chemical pregnancies" or "occult pregnancies." A landmark epidemiological study established that the incidence of

25

30

35

007080-5720E960

EPL was 22% in a population of healthy women attempting to conceive (Wilcox, A. J., et al., 1988). This investigation employed a very sensitive (0.01 ng/ml hCG) assay which detected only the intact hCG molecule with the unique beta subunit carboxyterminal peptide present.

There are multiple likely causes for EPL and clinical spontaneous abortion including genetic abnormality, immunological dysfunction, untreated infection or other unknown physiological problems. In addition, losses may be caused by failure of human chorionic gonadotropin (hCG) to induce adequate response at its target, the corpus luteum. This could result from inadequate hormonal potency. "Nicking" of the beta subunit in the loop 2 region of the molecule, specifically between residues 44-49, can reduce biopotency of hCG. Cleaved peptide bonds in this area of the molecule also exhibit reduced biopotency and reduced immunochemical recognition by monoclonal antibodies directed to the heterodimeric hormone (Cole, L. A., et al., 1991a; Cole, L. A., et al., 1991b; Puisieux, A., et al., 1990; Nishimura, R., et al., 1988; Nishimura, R. T., et al., 1989). Nicked forms of hCG were examined as possibly more prevalent in EPL situations and, at least in part responsible, for early pregnancy loss. Unfortunately many of the reports claiming that substantial concentrations of nicked hCG are produced during pregnancy, losses or successful pregnancies, are not accurate due to faulty assumptions regarding assay specificity. Carbohydrate-modified hCG can also exhibit either reduced or enhanced biopotency. It is known that if the hCG has much reduced sialic acid content and its carbohydrate chains terminate in galactose, much hCG would be removed by the liver receptor for such altered glycoproteins (Braun, J. R., et al., 1996; Kawasaki, T. and G. Ashwell, 1996). The circulating life-time of asialo hCG is reduced and its *in vivo* potency is thereby

007080-5720E950

low. Other carbohydrate changes also alter circulating half life; glycoproteins terminating in sulfate-N-acetyl galactosamine are also extracted by a specific liver receptor and have reduced circulating lifetime
5 (Baenziger, J. U., 1994; Fiete, D., et al., 1991).

At least two factors affect increased potency of hCG. First, it is known that a larger Stoke's radius will decrease clearance through the kidney glomerulus which generally clears proteins above an effective size of 70,000 very slowly. The effective size of urinary-isolated hCG is just at this borderline reduced clearance size. Generally, extra sugar content makes the hydrated radius of glycoproteins larger. It has been shown that by adding the hCG beta COOH-terminal peptide to hFSH or hLH, their circulating life-times greatly increased (Fares, F. A. et al., 1992; Matzuk, M. M., 1990). This addition was thought mostly due to the carbohydrate content of that peptide rather than simply the extra polypeptide size. Second, increased negative charge of a protein will prolong its circulating time because of decreased renal clearance (Chmielewski, C. 1992, Quadri, K. H., et al., 1994; Maack, T., et al., 1985). This increased negative charge can be due to extra sialic acid or other negative groups, including sulfate such as is present on hLH and on the pituitary form of hCG (Birken, S., et al., 1996b). Changes which affect signal transduction at the receptor may also affect biopotency of hCG. It is known that deglycosylated hCG has much reduced receptor potency (Ravindranath, N., et al., 1992; Sairam, M. R., and L. G., Jiang, 1992; Browne, E. S., et al., 1990; Sairam, M. R., 1989; Sairam, M. R., et al., 1988). Carbohydrate reduced forms of hCG also have reduced signal transduction (Amano, J., et al., 1990; Bahl, O. P., et al., 1995; Moyle, W. R., 1975).

001080" ST202560

According to the present invention EPL or recurrent
spontaneous abortion is not due to an abnormal hCG form
that has reduced potency, such as nicked hCG. Instead,
the present invention provides evidence that in
5 successful outcome pregnancies women usually produce
forms of hCG which are very highly potent in very early
pregnancy; the standard urinary reference preparations
of hCG are less potent forms of the hormone produced
later in pregnancy. The increased potency could be
10 caused by a combination of factors from circulating
half-life to increased receptor affinity or signal
transduction or all of the preceding. Since hCG is low
very early in pregnancy, it is logical to find a more
potent form of hCG on a molar basis to carry out its
15 function until production levels rise as the
trophoblastic cellular mass increases. The present
invention describes molecular and immunological tools
and methods including an antibody, B152, described
herein which recognizes the highly potent early
20 pregnancy associated molecular isoforms of hCG. The
determination of blood and urine profiles for the B152
hCG isoforms throughout healthy pregnancies can
delineate the pattern of isoforms in successful
pregnancies. These isoforms can be measured by
25 immunoassay alone, obviating the need to perform complex
isoelectric focusing studies or other separation
techniques. Additionally, the methods described herein
are applicable to large numbers of samples.

30

007080" ST20E560

Summary of the Invention

The present invention provides a method of predicting pregnancy outcome in a subject by determining the amount of an early pregnancy associated molecular isoform of hCG in a sample comprising: (a) contacting a sample with an antibody which specifically binds to the early pregnancy associated molecular isoform of hCG under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG; (b) measuring the amount of complexes formed, thereby determining the amount of the early pregnancy associated molecular isoform of hCG in the sample; and (c) comparing the amount early pregnancy associated molecular isoform of hCG in the sample determined in step (b) with the amount determined for temporally matched, normal pregnant subject(s) wherein the relative absence of the early pregnancy associated molecular isoform of hCG in the sample indicates a negative outcome of pregnancy for the subject.

The present invention further provides a method of predicting pregnancy outcome in a subject by determining the amount of an early pregnancy associated molecular isoform of hCG in a sample comprising: (a) contacting a capturing antibody which specifically binds to the early pregnancy associated molecular isoform of hCG with a solid matrix under conditions permitting binding of the antibody with the solid matrix; (b) contacting the bound matrix with the sample under conditions permitting binding of the antigen present in the sample with the capturing antibody; (c) separating the bound matrix and the sample; (d) contacting the separated bound matrix with a detecting antibody which specifically binds to hCG under conditions permitting binding of antibody and antigen in the sample; (e) measuring the amount of bound antibody on the bound matrix, thereby determining the amount of early pregnancy associated molecular isoform

001080 " ST 202560

of hCG in the sample; and (f) comparing the amount early pregnancy associated molecular isoform of hCG in the sample determined in step (e) with the amount determined for temporally matched, normal pregnant subject(s), wherein amounts of the early pregnancy associated molecular isoform of hCG in the sample similar to amounts of early pregnancy associated molecular isoform of hCG in temporally matched pregnant samples indicates a positive outcome, amounts of early pregnancy associated molecular isoform of hCG in the sample similar to amounts of early pregnancy associated molecular isoform of hCG in the non-pregnant samples indicates a negative outcome of pregnancy for the subject.

In addition, the present invention provides a method for determining the amount of early pregnancy associated molecular isoforms of in a sample comprising: (a) contacting the sample with an antibody which specifically binds to an early pregnancy associated molecular isoform of hCG under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG; and (b) determining the amount of complexes formed thereby determining the amount of early pregnancy associated molecular isoform of hCG in the sample.

Further, the present invention provides a diagnostic kit for determining the amount of early pregnancy associated hCG is a sample comprising: (a) an antibody which specifically binds to an early pregnancy associated molecular isoform; (b) a solid matrix to which the antibody is bound; and (c) reagents permitting the formation of a complex between the antibody and a sample.

The present invention additionally provides an antibody

007030" ST20E960

which specifically binds to an early pregnancy associated molecular isoform of human chorionic gonadotropin.

5 Further, the present invention provides a method for
detecting non-trophoblast malignancy in a sample
comprising: (a) contacting a sample with an antibody
which specifically binds to the early pregnancy
associated molecular isoform of hCG under conditions
10 permitting formation of a complex between the antibody
and the early pregnancy associated molecular isoform of
hCG; (b) contacting the sample with a second antibody
which specifically binds to intact non-nicked hCG
without substantially cross-reacting with said antibody
15 under conditions permitting formation of a complex
between the antibody and the early pregnancy associated
molecular isoform of hCG; (c) measuring the amount of
complexes formed, thereby determining the amount of the
early pregnancy associated molecular isoform of hCG in
20 the sample; and (d) comparing the amount of early
pregnancy associated molecular isoform of hCG in the
sample determined in step (b) with the amount of early
pregnancy associated molecular isoform of hCG in the
sample determined in step (c), wherein a positive
25 detection of early pregnancy associated molecular
isoform detected in step (b) and a relative absence of
the early pregnancy associated molecular isoform of hCG
detected in step (c) indicates the presence of non-
trophoblast malignancy in the sample.

30 Finally, the present invention provides a method for
detecting gestational trophoblast disease in a sample
from a subject comprising (a) contacting a sample with
an antibody which specifically binds to the early
pregnancy associated molecular isoform of hCG under
35 conditions permitting formation of a complex between the
antibody and the early pregnancy associated molecular

SECRET 090950

isoform of hCG; (b) contacting the sample with a second antibody which specifically binds to intact non-nicked hCG without substantially cross-reacting with said antibody under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG; (c) measuring the amount of complexes formed, thereby determining the amount of the early pregnancy associated molecular isoform of hCG in the sample due to binding with the first antibody, and late pregnancy associated molecular isoform of hCG in the sample due to binding with the second antibody; (d) determining the ratio of early pregnancy associated molecular isoform of hCG to late pregnancy associated molecular isoform of hCG in the subject; and (e) comparing the ratio of early pregnancy associated molecular isoform of hCG to late pregnancy associated molecular isoform of hCG in the sample determined in step (c) over time, wherein a continuing high ratio of early pregnancy associated molecular isoform of hCG to late pregnancy associated molecular isoform of hCG in the sample determined in step (c) indicates the presence of gestational trophoblast disease in the subject.

5

10

15

20

25

001000 ST20E960

Brief Description of the Figures

Figure 1.

Bioassay for forms of hCG. This is data from recombinant CHO cells expressing the LH/CG receptor. The response factor is cAMP production. The x-axis is dose of one of four calibrated, pure hormones as described on graph legends. Expressed hCG has no nicks; choriocarcinoma hCG (C5) is 100% nicked; CR 127 was purified into a nick-free (intact) and nick-enriched fraction as shown.

5
10

Figure 2.

Incidence (Panel A) and expression level (Panel B) of hCG-related molecules in the positive samples for each of the analyses measured (In early normal pregnancy, n=214; EPL cycles, n=49; and negative cycles, n=297).

15

Figure 3.

Binding curves for three hCG types in the B152-B207* assay (upper panel) and the B109-B108* assay (lower panel).

20

Figure 4.

Ratio of hCG isoforms measured by the B152-B207* and B109-B108* assays in normal pregnancy urine (n=103) at different gestational ages. (Regression curve and 95% confidence intervals are shown, $r^2=0.79$). An inflection point in the curve occurs at approximately 29 weeks.

25

Figure 5.

Box plot of the B152/B109 ratio for pregnancy matched serum/urine at 5-6 weeks of gestational age (n=12); or at 36-39 weeks of gestational age (n=11) and in JAR cell supernatant. Box extends to the 25th and 75th percentile. The upper and lower symbols indicate the 90th and 10th percentile respectively. A solid line inside the box marks the value of the 50th percentile.

30

35

007030*5720E960

Figure 6.

Ratio of hCG isoforms measured by the B152-B207* and B109-B108* assays in the urine of IVF patients (n=65). (Regression curve and 95% confidence intervals are shown, $r^2=0.59$).

5

Figure 7.

Immunoassay profiles of fractions from Superose 12 column chromatography of a pooled urine concentrate from pregnant women.

10

007080"ST20E960

Detailed Description of the Invention

5 A method of predicting pregnancy outcome in a subject by
determining the amount of an early pregnancy associated
molecular isoform of hCG in a sample comprising: (a)
10 contacting a sample with an antibody which specifically
binds to the early pregnancy associated molecular
isoform of hCG under conditions permitting formation of
a complex between the antibody and the early pregnancy
associated molecular isoform of hCG; (b) measuring the
15 amount of complexes formed, thereby determining the
amount of the early pregnancy associated molecular
isoform of hCG in the sample; and (c) comparing the
amount early pregnancy associated molecular isoform of
20 hCG in the sample determined in step (b) with either (i)
the amount determined for temporally matched, normal
pregnant subject(s) or (ii) the amount determined for
non-pregnant subject(s), wherein the relative absence of
the early pregnancy associated molecular isoform of hCG
in the sample indicates a negative outcome of pregnancy
for the subject. In an embodiment of the present
invention, the antibody is B152. Another embodiment of
this invention is the early pregnancy associated
molecular isoform of hCG.

25 The hybridoma producing the B152 monoclonal antibody was
deposited on February 3, 1998 with the American Type
Culture Collection (ATCC), 12301 Parklawn Drive,
Rockville, Maryland 20852, U.S.A. under the provisions
30 of the Budapest Treaty for the International Recognition
of the Deposit of Microorganism for the Purposes of
Patent Procedure. The hybridoma, was accorded ATCC
Accession Number HB-12467.

35 According to one embodiment of this invention, step (a)
further comprises a second antibody which specifically
binds to hCG without substantially cross-reacting with

007080" ST00E960

5 said antibody under conditions permitting formation of
a complex between the antibody and the early pregnancy
associated molecular isoform of hCG. In an embodiment
of this invention, the second detection antibody is
10 B207. According to another embodiment of this invention,
step (a) further comprises a second assay antibody B109
which specifically binds to intact non-nicked hCG
without substantially cross-reacting with said antibody
under conditions permitting formation of a complex
15 between the antibody and the early pregnancy associated
molecular isoform of hCG. In an embodiment of this
invention, the detection antibody is B108. In an
embodiment of this invention, step (c) comprises
comparing the amount of the early pregnancy associated
20 molecular isoform of hCG determined in step (b) for
B152-B207 assay with the amount determined in step (b)
for the B109-B108 assay wherein a high ratio of amounts
determined for said antibody relative to the second
antibody indicates a positive outcome of pregnancy for
the subject, a low ratio indicates a negative outcome of
pregnancy for the subject.

In yet another embodiment of this invention, step (c)
comprises comparing the amount early pregnancy
25 associated molecular isoform of hCG in the sample
determined in step (b) with the amount determined for
temporally matched, normal pregnant subject(s), wherein
amounts of the early pregnancy associated molecular
isoform of hCG in the sample similar to amounts of early
30 pregnancy associated molecular isoform of hCG in
temporally matched pregnant samples indicates a positive
outcome, amounts of early pregnancy associated molecular
isoform of hCG in the sample similar to amounts of early
pregnancy associated molecular isoform of hCG in the
35 non-pregnant samples indicates a negative outcome of
pregnancy for the subject.

001030"51202960

This invention also provides a method of predicting the likelihood of a negative pregnancy outcome in a female subject comprising: (a) contacting a sample from the subject with a capture antibody which specifically binds to an early pregnancy associated molecular isoform of hCG under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG; (b) contacting any complex formed in step (a) with a labelled detection antibody under conditions permitting binding to the complex the capture antibody and the hCG isoform; (c) measuring the amount of labeled detection antibody bound to the complex so as to thereby determine the amount of the early pregnancy associated molecular isoform of hCG in the sample; and (d) comparing the amount early pregnancy associated molecular isoform of hCG in the sample determined in step (b) with the amount determined for a normal pregnant subject, wherein the relative absence of the early pregnancy associated molecular isoform of hCG in the sample indicates a negative outcome of pregnancy for the subject.

According to an embodiment of this invention, the sample is a urinary sample or a blood sample. In one embodiment of this invention, the sample is an aggregate sample taken from at least one day. In another embodiment, sample may be taken from at least two consecutive days and in a further embodiment, the sample is taken in three days. In an embodiment of this invention, the sample is a spot urine sample, a first morning void urine sample, or an aggregate sample of the first morning void urine samples for at least two consecutive days. In one embodiment of this invention, the antibody is labeled with a detectable marker. In an embodiment of this invention, the detectable marker is a radioactive isotope, enzyme, dye, magnetic bead, or biotin. In a preferred embodiment, the radioactive

001030 "STRE960

isotope is I¹²⁵.

5 The present invention further provides a method of predicting pregnancy outcome in a subject by determining the amount of an early pregnancy associated molecular isoform of hCG in a sample comprising: (a) contacting a capturing antibody which specifically binds to the early pregnancy associated molecular isoform of hCG with a solid matrix under conditions permitting binding of the antibody with the solid matrix; (b) contacting the bound matrix with the sample under conditions permitting binding of the antigen present in the sample with the capturing antibody; (c) separating the bound matrix and the sample; (d) contacting the separated bound matrix with a detecting antibody which specifically binds to hCG under conditions permitting binding of antibody and antigen in the sample; (e) measuring the amount of bound antibody on the bound matrix, thereby determining the amount of early pregnancy associated molecular isoform of hCG in the sample; and (f) comparing the amount early pregnancy associated molecular isoform of hCG in the sample determined in step (e) with the amount determined for temporally matched, normal pregnant subject(s), wherein amounts of the early pregnancy associated molecular isoform of hCG in the sample similar to amounts of early pregnancy associated molecular isoform of hCG in temporally matched pregnant samples indicates a positive outcome, amounts of early pregnancy associated molecular isoform of hCG in the sample similar to amounts of early pregnancy associated molecular isoform of hCG in the non-pregnant samples indicates a negative outcome of pregnancy for the subject.

35 An embodiment of this invention further comprises (a) removing of the sample from the matrix; and (b) washing the bound matrix with an appropriate buffer. In one

OFFICE "STENO" 0000

embodiment of this invention, the capturing antibody is B152. In one embodiment of this invention, the detecting antibody is B207. In an embodiment of this invention, step (a) further comprises a second capturing antibody which specifically binds to intact non-nicked hCG without substantially cross-reacting with said antibody under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG. According to an embodiment of this invention, the second capturing antibody is B109 and the second detection antibody is B108. In an embodiment of this invention, step (d) further comprises a second detecting antibody which specifically binds to hCG without substantially cross-reacting with said antibody under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG. In an embodiment of this invention, step (f) comprises comparing the amount of the early pregnancy associated molecular isoform of hCG determined in step (e) for said antibody with the amount determined in step (b) for the second antibody, wherein a high ratio of amounts determined for said antibody relative to the second antibody indicates a positive outcome of pregnancy for the subject, a low ratio indicates a negative outcome of pregnancy for the subject.

According to an embodiment of this invention, the sample is a urinary sample or a blood sample. In one embodiment of this invention, the sample is an aggregate sample taken from at least two consecutive days. In an embodiment of this invention, the sample is a spot urine sample, a first morning void urine sample, or an aggregate sample of the first morning void urine samples for at least two consecutive days. In one embodiment of this invention, the antibody is labeled with a detectable marker. In an embodiment of this invention,

00100 * 5T2E2960

the detectable marker is a radioactive isotope, enzyme, dye, magnetic bead, or biotin. In a preferred embodiment, the radioactive isotope is I¹²⁵.

5 In addition, the present invention provides a method for determining the amount of early pregnancy associated molecular isoforms of in a sample comprising: (a) contacting the sample with an antibody which specifically binds to an early pregnancy associated
10 molecular isoform of hCG under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG; and (b) determining the amount of complexes formed thereby determining the amount of early pregnancy associated
15 molecular isoform of hCG in the sample.

According to an embodiment of this invention, the antibody specifically binds a region of the early pregnancy associated molecular isoform of hCG comprising
20 a carbohydrate moiety. In one embodiment of this invention the antibody is produced by a hybridoma cell line. In one embodiment of this invention the antibody is B152.

25 Further, the present invention provides a diagnostic kit for determining the amount of early pregnancy associated hCG is a sample comprising: (a) an antibody which specifically binds to an early pregnancy associated molecular isoform; (b) a solid matrix to which the
30 antibody is bound; and (c) reagents permitting the formation of a complex between the antibody and a sample. In an embodiment of this invention, the antibody is B109 or B152. An embodiment of this invention further comprises control sample(s) normal
35 pregnant sample(s), nonpregnant sample(s), or male sample(s). The kit may also contain detection antibodies such as B207 or B108.

0301080" 57202960

According to an embodiment of this invention, the sample is a urinary sample or a blood sample. In one embodiment of this invention, the sample is an aggregate sample taken from at least one or may be two or three consecutive days. In an embodiment of this invention, 5 the sample is a spot urine sample, a first morning void urine sample, or an aggregate sample of the first morning void urine samples for at least one, or may be two or three consecutive days. In one embodiment of this invention, the antibody is labeled with a detectable 10 marker. In an embodiment of this invention, the detectable marker is a radioactive isotope, enzyme, dye, magnetic bead, or biotin. In a preferred embodiment, the radioactive isotope is I^{125} .

15 The present invention additionally provides an antibody which specifically binds to an early pregnancy associated molecular isoform of human chorionic gonadotropin.

20 In an embodiment of this invention, the antibody specifically binds to a region of the early pregnancy associated molecular isoform of human chorionic gonadotropin comprising a carbohydrate moiety.

25 According to one embodiment of this invention, the monoclonal antibody is B152. In an embodiment of this invention, a hybridoma cell (ATCC Accession No. HB-12467) is provided capable of producing monoclonal antibody B152. Another embodiment of this invention is 30 the early pregnancy associated molecular isoform of hCG recognized by the B152 monoclonal antibody.

Further, the present invention provides a method for 35 detecting non-trophoblast malignancy in a sample comprising: (a) contacting a sample with an antibody which specifically binds to the early pregnancy associated molecular isoform of hCG under conditions

0933015 080100
001000 ST200960

001000 "ST20E60

5 permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG; (b) contacting the sample with a second antibody which specifically binds to intact non-nicked hCG
10 without substantially cross-reacting with said antibody under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG; (c) measuring the amount of complexes formed, thereby determining the amount of the early pregnancy associated molecular isoform of hCG in the sample; and (d) comparing the amount of early pregnancy associated molecular isoform of hCG in the sample determined in step (b) with the amount of early pregnancy associated molecular isoform of hCG in the sample determined in step (c), wherein a positive detection of early pregnancy associated molecular isoform detected in step (b) and a relative absence of the early pregnancy associated molecular isoform of hCG detected in step (c) indicates the presence of non-trophoblast malignancy in the sample.
20

25 According to an embodiment of this invention, the antibody is B152 or B109. In an embodiment of this invention, the detection antibody is B207 for B152 assay, B108 for B109 assay. In an embodiment of this invention, the non-trophoblast malignancy is ovarian malignancy or prostate malignancy.

30 According to an embodiment of this invention, the sample is a urinary sample or a blood sample. In one embodiment of this invention, the sample is an aggregate sample taken from at least two consecutive days. In an embodiment of this invention, the sample is a spot urine sample, a first morning void urine sample, or an aggregate sample of the first morning void urine samples for at least two consecutive days. In one embodiment of this invention, the antibody is labeled with a
35

detectable marker. In an embodiment of this invention, the detectable marker is a radioactive isotope, enzyme, dye, magnetic bead, or biotin. In a preferred embodiment, the radioactive isotope is I¹²⁵.

5

Finally, the present invention provides a method for detecting gestational trophoblast disease in a sample from a subject comprising (a) contacting a sample with an antibody which specifically binds to the early pregnancy associated molecular isoform of hCG under
10 conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG; (b) contacting the sample with a second antibody which specifically binds to intact non-nicked hCG without substantially cross-reacting with
15 said antibody under conditions permitting formation of a complex between the antibody and the early pregnancy associated molecular isoform of hCG; (c) measuring the amount of complexes formed, thereby determining the amount of the early pregnancy associated molecular
20 isoform of hCG in the sample due to binding with the first antibody, and late pregnancy associated molecular isoform of hCG in the sample due to binding with the second antibody; (d) determining the ratio of early pregnancy associated molecular isoform of hCG to late pregnancy associated molecular isoform of hCG in the
25 subject; and (e) comparing the ratio of early pregnancy associated molecular isoform of hCG to late pregnancy associated molecular isoform of hCG in the sample determined in step (c) over time, wherein a continuing high ratio of early pregnancy associated molecular isoform of hCG to late pregnancy associated molecular isoform of hCG in the sample determined in step (c) indicates the presence of gestational trophoblast
30 disease in the subject.
35

In an embodiment of this invention, the antibody is B152

001030 " 5120950

or B109. In another embodiment of this invention, the detection antibody is B108 for B109 assay, B207 for B152 assay. In an embodiment of the present invention, the gestational trophoblast disease is choriocarcinoma or hydatidiform mole.

According to an embodiment of this invention, the sample is a urinary sample or a blood sample. In one embodiment of this invention, the sample is an aggregate sample taken from at least two consecutive days. In an embodiment of this invention, the sample is a spot urine sample, a first morning void urine sample, or an aggregate sample of the first morning void urine samples for at least two consecutive days. In one embodiment of this invention, the detection antibody B207 or B108 is labeled with a detectable marker. In an embodiment of this invention, the detectable marker is a radioactive isotope, enzyme, dye, magnetic bead, or biotin. In a preferred embodiment, the radioactive isotope is I¹²⁵.

As described herein below, unexpected isoforms of hCG are produced during normal early pregnancy. Using an *in vitro* bioassay, it appears that these isoforms have enhanced potency for signal transduction. These isoforms can be measured using the novel sensitive, immunoassay described herein. This can help predict pregnancy outcome where one cause of early pregnancy loss is failure to produce the isoform of hCG of higher potency produced by successful pregnancies. This enables physicians to intervene to sustain a failing pregnancy. Identification of the nature of the hCG isoform required might provide the proper reagent needed to sustain pregnancy.

New antibodies for measurement of nicked forms of hCG described herein below were developed based on the hypothesis that forms of hCG, which have greatly reduced

09830801500000

bioactivity, contribute to early pregnancy loss (EPL), due at least in part to diminished biopotency. Evidence was found that the hCG that appears in EPL patients displays reduced biological activity. However, it was
5 determined that the cause of the reduced bioactivity is not the presence of nicked hCG in EPL patients. Instead, the hypothesis is that patients that carry pregnancies forward produce an isoform of hCG with enhanced bioactivity. The instant invention describes
10 a unique immunochemical assay to measure this unexpected and previously un-characterized isoform of early pregnancy hCG directly in clinical samples of blood and urine. One of the antibodies developed reacted against a nicked form of hCG isolated from a choriocarcinoma
15 patient, was not specific for a nicked form of hCG but appeared to discriminate among carbohydrate variants of hCG. This antibody, designated B152, appears to preferentially bind hCG forms from choriocarcinoma patients. In studying the content of hCG isoforms
20 during pregnancy, the unique and unexpected observation was made that B152 in the first four weeks of pregnancy measured much higher quantities of an isoform of hCG as compared to the standard hCG isoforms measured by the usual heterodimeric hCG assays exemplified by a
25 previously described B109 based assay. In fact, in early pregnancy (days 9,10,11 postovulation) B152 measured as much as 20-fold more hCG, than did another monoclonal antibody, B109. Later in pregnancy, the B152 isoform declines and is lower in third trimester
30 pregnancy urine than the standard isoforms measured by B109. A further striking observation was that in very early pregnancy, a high B152/B109 ratio correlates with a successful pregnancy outcome while a low ratio correlated with pregnancy loss. This discovery is
35 important as the potentially overlooked isoforms of hCG described herein during pregnancy may be predictors of successful pregnancy outcome. Such an assay has wide

09530315 00100

medical applications and provides a clinician with opportunity to intervene very early in pregnancy if the assay indicated that the pregnancy appeared troubled.

5 An antibody, designated B152, produced by the hybridoma cell accorded ATCC Accession number HB-12467 generated against a nicked form of hCG isolated from a choriocarcinoma patient, but not specific for nicked isoform hCG is able to discriminate among carbohydrate variants of hCG. B152 is specific for an early pregnancy associated molecular isoform of hCG. which in 10 the first four weeks of pregnancy is measured at much higher quantities than the hCG standard isoforms measured by the usual heterodimeric hCG assays exemplified by a previously described B109 based assay. 15 Later in pregnancy, the B152 isoform declines and is lower in third trimester pregnancy urine than the standard isoforms measured by B109.

20 This invention is illustrated in the Experimental Details section which follows. These sections are set forth to aid in an understanding of the invention but are not intended to, and should not be construed to, limit in any way the invention as set forth in the 25 claims which follow thereafter.

09290215 080100

EXPERIMENTAL DETAILS

Example 1: **Analysis of Molecular Isoforms of hCG in
Early Pregnancy and Early Pregnancy Loss**

5

Introduction

Almost all investigations of the incidence of early pregnancy loss (EPL), either in normal populations or in populations at risk as a consequence of exposure to putative reproductive toxins (Hakim, R. B., et al., 1995; Lasley, B. L., et al., 1995) use assays for heterodimeric, non-nicked hCG or combination assays which additionally include free beta subunit and beta core fragment of hCG. One concern about the forms of hCG to include in the measurement in EPL was heightened with respect to the nicking phenomenon described above. Because nicked hCG molecules are not measured by the antibodies employed in most EPL studies, the incidence of EPL is presumably underestimated by an amount proportional to the extent of nicking in the urinary molecule. Another concern of significant importance was a determination of the nature of the "hCG like" immunoreactivity in the urine in the periovulatory surge of the menstrual cycle (O'Connor J., et al., 1995). Recent reports have confirmed the existence of and documented the structure of a sulfated form of hCG produced in the pituitary (Birken, S., et al., 1996b). There is a pulsatile secretion of hCG in both men and non-pregnant women. (Odell, W. D.; Griffin, J., 1989 and Odell, W. D.; Griffin, J., 1987). The presence of a non-pregnancy associated form of sulfated hCG of pituitary origin, peaking at ovulation and perhaps persisting into the luteal phase, could potentially interfere with the accurate estimation of EPL.

35

Unappreciated isoforms of hCG in blood and urine very early in pregnancy may be more potent *in vivo* than the forms of hCG produced later in pregnancy. The absence

001030"5F20E560

of such isoforms may be one cause of early pregnancy
loss. A sensitive and specific immunoassay system was
designed and made to measure unique early pregnancy
associated molecular isoforms (EPMI) of hCG. These
5 isoforms, likely to differ by carbohydrate composition,
are predictive of a successful pregnancy outcome. When
these early pregnancy associated molecular isoforms of
hCG are absent or present in low concentration, the
pregnancy may be lost very early and be observed as only
10 a "chemical" pregnancy. These hCG isoforms may resemble
the forms of hCG produced in some choriocarcinoma
patients from which the immunogen used to produce
monoclonal antibody B152 was derived as described herein
below. The isoforms resemble those from trophoblastic
15 disease not in terms of nicking or intact peptide chains
but likely in carbohydrate content. The present
invention describes that the molar ratio of B152 to B109
epitopes are predictive of a successful pregnancy or a
loss. Three categories of pregnant patients were
20 analyzed: (a) normal pregnant women, (b) women who
experience recurrent abortions, (c) women undergoing
embryo implantation.

It is possible to determine the hCG isoforms present in
25 the blood and urine of women who have a history of
recurrent spontaneous abortion and a similar analysis of
women undergoing embryo implantation. The combined EPL
and spontaneous abortion rate in healthy populations is
31%. Subjects who experience three consecutive recurrent
30 spontaneous abortions have a 32% risk of sustaining
another (Hill, J. A.; Anderson, D. J., 1990). In in
vitro fertilization IVF pregnancy, the loss rate is 70%
with non-donor sperm and 50% when donor sperm is used.
Delineation of pregnancies with a negative outcome from
35 pregnancies with a positive outcome can be based on
differences in the concentrations of EPMI hCG isoforms
(i.e. as differences in the B152/B109 ratio in

0010159130

patients). In addition, specimens from gestational trophoblastic disease (GTD) can be used to discriminate between GTD and normal pregnancy.

Results

5 In vitro bioassay for hLH/hCG

An hCG bioassay was constructed employing CHO cells expressing functional human LH/CG receptor. Figure 1 illustrates the differences in vitro in biological activity between nicked and non-nicked hCG as measured by this assay. This system, has been used to evaluate the activity of pituitary and placental hCG (Birken, S., et al., 1996b). Preparations of hCG were tested for nicked and non-nicked molecular isoforms of hCG in a second recombinant bioassay system (Ho, H-H., et al., 1997). Similar results were obtained in both systems.

Normal pregnancy values compared with EPL values.

Figure 2 indicated that nicked hCG is not a significant molar constituent of either early pregnancy or EPL. Data indicated that biological activity is not correlated with nicked hCG, but is instead ascribed to a form of hCG recognized by the B152 monoclonal antibody -- an early pregnancy associated molecular isoform of hCG (EPMI hCG). It has been established that there is diminished hCG bioactivity associated with EPL as compared to early normal pregnancy (Ho, H-H., et al., 1997). Thus, diminished hCG biological activity is a factor in EPL as a consequence of a heretofore unappreciated isoform of hCG - an early pregnancy associated molecular isoform of hCG.

hCG Urinary Analytes. Metabolites of hCG and hLH were studied in a variety of states (Birken, S., et al., 1996a). One study indicated a 31% pregnancy loss (Zinaman, MJ, et al., 1996) while another indicated a 17.4% rate of early pregnancy loss based on hCG assays (Elish, N. J., et al., 1996). It is known that hCG and

00630215-080100

hCG beta core can be readily transferred from the uterus to the circulation even in the absence of implantation (Chang, P. L., 1997). The molecular spectrum of hCG urinary analytes in EPL cycles, normal conceptive cycles 5 and non-conceptive cycles has been evaluated. The study design and demographics of the investigation have been described (Ellish, N. J., et al., 1996).

Briefly, three urine specimens per cycle, corresponding 10 to days 9,10, 11, post calculated day of ovulation were collected and analyzed in a screening assay (the "combo") which simultaneously detects intact, non-nicked hCG, hCG free beta subunit, and hCG beta core fragment. Individual determinations for each of these analytes, 15 as well as for nicked hCG, and the form of intact hCG detected by monoclonal antibody B152 (EPMI hCG) were performed on these specimens. In addition, since the concentration of luteal phase hLH urinary analytes is a concern because of cross-reaction in hCG assays, levels 20 of intact hLH, hLH free beta subunit and hLH beta core fragment were determined in the normal pregnancy cycles and the non-conceptive cycles. Table I summarizes the characteristics of immunometric assays employed.

25

000000000000000000000000

TABLE 1. Assay format and specificity

<i>Assay format</i>	<i>Primary analyte</i>	<i>% cross-reactivity with related analytes</i>	<i>Intra-assay cv, %</i>	<i>Inter-assay cv, %</i>
B109-B108*	intact non-nicked hCG	<1% ^b	6	12
B201-C104*	hCG free beta subunit (non-nicked+nicked)	1% hCG; 10% hCG nicked (pregnancy); <1% ^b	6	12
B210-B108*	hCG beta core fragment	2% hLH beta core fragment; <1% ^b	5	7
B151-B207*	hCG nicked	10% hCG nicked free beta subunit; 12% hCG non-nicked; 2% hCG free beta subunit; 2% hLH; 5% hLH free beta subunit; <1% ^b	5	15
B152-B207*	choriocarcinoma hCG (C5) and choriocarcinoma hCG free beta subunit	100% hCG nicked (C5); 190% hCG free beta nicked (from C5); 10% hCG nicked (pregnancy); 5% hCG free beta nicked (pregnancy); 7% hCG (pregnancy); 6% hCG free beta subunit; <1% ^b	6	13
B406-A201*	hLH	<1% ^a	4	10
B505-B503*	hLH beta core fragment	<1% ^a	9	9
B408-B409*	hLH free beta subunit	29% hLH; <1% ^a	7	11

^a-(if not indicated) hLH, free beta hLH, hLH beta core fragment, hCG, free beta hCG, hCG beta core fragment;

^b-(if not indicated) the same as ^(a) plus nicked hCG and nicked free beta hCG (pregnancy).

001080"5F20E960

The results indicate that nicked hCG does not constitute
 a significant mole fraction of urinary hCG
 immunoreactivity in either EPL or early normal
 pregnancy. In addition, there is a substantial
 5 excretion of hCG free beta subunit in some subjects in
 both pregnancy and EPL. Further, both EPL and normal
 pregnancy cycles variably express all of the measured
 analytes. Although both the incidence and level of
 expression are different between EPL's and normal
 10 pregnancy, there is no hCG related analyte unique to
 either state. There was, however, a clear difference
 between the hLH associated analytes in the control
 population (non-conceptive cycles) and the normal
 pregnancy group. Virtually all of the non-pregnancy
 15 cycles expressed hLH free beta subunit and hLH beta core
 fragment while only a third of the conceptive cycles had
 detectable levels of either analyte. Intact hLH proved
 to be a minor constituent of the hLH profile in both
 groups.

20 These findings demonstrate both the necessity of
 measuring hCG beta core fragment in the detection of
 EPL, and also of making sure that the hCG beta core
 assay does not cross-react with beta core hLH, which is
 25 demonstrated to be present in that part of the luteal
 phase where EPL measurements are performed. The data is
 summarized in Figure 2.

30 Statistical analysis was performed after transformation
 of analyte values to mole fractions so as to produce a
 more useful analysis due to the wide excursion of hCG
 analyte values among groups. The mole fraction data
 were evaluated by discriminant analysis and by a mixed
 effects model incorporating LMP. The discriminant
 35 analysis was performed both with and without "outliers"
 (defined as values greater than two standard deviation
 from the mean) removed. Both approaches produced

001000"5F20E960

recognition is quite different. The B152-B207* assay is more sensitive to C5, which is to be expected because B152 antibody was developed and selected on the basis of higher affinity to C5. Urinary non-nicked hCG is purified from the CR127 preparation of pooled normal pregnancy hCG. Conversely C5 is recognized with lower affinity by the B109-B108* assay, which has primary specificity for the hCG isoforms of later pregnancy.

We have developed a method to directly profile changes of hCG isoforms in serum or urine throughout pregnancy. Two IRMAs for hCG are employed, each based on monoclonal antibodies to different hCG epitopes. The B109-B108* assay is a commonly used intact hCG assay to the heterodimeric-dependent epitope. A new assay, B152-B207*, is most likely sensitive to the carbohydrate portion of hCG carboxyterminal peptide. The same standard non-nicked hCG was used in both assays. Non-nicked hCG was employed since the B109 assay reacts poorly with nicked forms of hCG while the B152 assay does not discriminate between nicked and non-nicked forms of the hormone. The B152 assay detected with greatly enhanced sensitivity hCG isoforms which appear earlier in pregnancy than isoforms measured by the B109 assay (O'Connor et al. 1998). Prior to development of the new immunometric assay system described in this report, it was not possible to readily discern the changes in hCG isoforms from very early pregnancy to mid pregnancy. The only available procedure for examining these changes was isoelectric focusing of every patient specimen followed by immunoassay of every focused fraction (Berger et al. 1993; Ulloa-Aguirre et al. 1990). The IEF pattern reflects the heterogeneity of the charged sugar, sialic acid which varies with the multi-antennary structures of the carbohydrate moieties in which sialic acid is the terminal sugar. Although we do not yet know the precise nature of the isoform epitopes

09630215-08100

being measured, the evidence for carbohydrate discrimination is based upon the hyperglycosylated structure of the immunogen, C5, used to develop the B152 monoclonal antibody and the antibody's reactivity with the hCG isoforms found in the JAR choriocarcinoma cell line. C5 hCG was isolated from a choriocarcinoma patient and has been thoroughly characterized as to its protein and carbohydrate content and structure (Elliott et al. 1997). It has been shown that C5 (and hCG from other choriocarcinoma subjects) differ in the protein moiety mainly by the presence of an increased number of nicked sites and by increased glycosylation relative to the hCG of normal pregnancy. In comparison with the hCG of normal pregnancy, choriocarcinoma derived hCG has increased fucosylation of the N-linked biantennary oligosaccharides in the beta subunit. In addition, the O-linked oligosaccharides in preparation C5 (a form of hCG produced from a single patient with choriocarcinoma) has a 100% tetrasaccharide core on the COOH-terminal region of the beta subunit. Normal mid pregnancy hCG has only 10-20% of this structure (Elliott et al. 1997). These observations, plus our own determination that the hCG synthesized by the JAR choriocarcinoma cell line provides a B152/B109 isoform ratio similar to that observed in early pregnancy, leads us to the conclusion that in very early pregnancy, the developing trophoblast secretes an isoform of hCG which resembles that produced in choriocarcinoma.

We have also tested recognition of pituitary hCG since its N-Asn carbohydrates differ somewhat from those of placental hCG, bearing a closer resemblance to those of hLH which have both sialic acid and sulfate groups (Birken et al. 1996). The carbohydrate structure of the COOH-terminal portion of pituitary hCG is not yet known. Since B152 did not recognize any substantial differences between pituitary and placental hCG (Figure

09530215 100100

3), differences in N-Asn recognition are unlikely. In terms of the COOH-terminal carbohydrates, it appears that pituitary and placental hCG (mid-pregnancy isoforms) may be similar, assuming the O-linked carbohydrate on the C5 antigen is part of the epitope of B152.

10 **Example 2: B152/B109 Ratio Predicts Pregnancy**
Outcome

The B152/B109 ratio measured in urine samples throughout the pregnancy

15 The relative concentrations of hCG isoforms in 103 normal pregnancy urine samples (5-39 weeks post last menstrual period - LMP) were determined by two immunometric assays (B152-B207* and B109-B108*). Both because of the wide range of hCG concentrations in
20 different samples, even at the same gestational age, and because neither of the assays is totally specific for the two (or more) families of hCG isoforms present, we find that presenting the data as a ratio of the observed two isoform groups more clearly delineates the change in
25 isoform content as pregnancy progresses. This calculated ratio is shown in Figure 4. In weeks 5-8 of pregnancy, the ratio of B152/B109 isoforms ranged between 6.2 and 1.3, indicating a predominance of the B152 isoform(s) in early pregnancy. During the 10 to 12
30 week period, the ratio ranged from 1 - 0.2, indicating that an inversion in hCG isoform content is occurring as pregnancy progresses. This decline in the ratio continues, ranging from 0.54 - 0.08 in the 15 - 18 week period and reaching an inflection point at 29 weeks. At
35 that time, the ratio reached a value of around 0.06 after which the ratio displayed a rise to a range of 0.2 - 0.07 in the 37 - 39.5 weeks of gestation time period.

09630215-080100

Statistical analysis involved fitting the log transformed ratio data to second and third order polynomial regression models. Since the third order term was not significant (likelihood ratio $\chi^2(1)=1.32$, $P=0.25$), the second order model was used ($r^2=0.793$). The log B152/B109 ratio reached an inflection point at LMP=29 weeks, based on this model.

The B152-B207* values reflect a measurement of the B152 isoform in terms of later pregnancy hCG equivalents, not in absolute quantities. It must be emphasized that the "absolute" concentrations measure in the B152 assay cannot be compared with the results of the B109 assay on an equimolar basis since the potency of the hyperglycosylated isoform is much higher in the B152 assay vis-à-vis the standard, i.e. normal later first trimester pregnancy hCG. The actual molar values of this isoform are on the order of tenfold less than those recorded in the assay. For this reason we have chosen not to analyze absolute molar quantities of the two analytes but only the ratio of the two measurements.

Even in normal pregnancy, the hCG values obtained vary widely according to the characteristics of the immunological reagents employed (Cole and Kardana, 1992; Cole et al. 1993). We hypothesize that the two assays described in this report primarily detect hCG isoforms at opposing ends of this spectrum, each primarily recognizing a subset of closely related molecules in the continuum of early to later pregnancy hCG molecular forms.

We have retained the use of normal pregnancy hCG as the standard in B152-B207* assay, despite its decreased affinity in this antibody configuration. The reasons for this include the limited and unrenewable supply of C5 (which was isolated from the urine of a single

09530215-080100

patient) and the variability in data which would result from investigations using different standards. The consequences of this choice are that the early pregnancy hCG isoforms have markedly increased immunopotency over
5 that of normal pregnancy and hence their molar quantities are overestimated in this assay. We use this difference in affinity to our advantage by employing a ratio of the molar results of two assays (B152 and B109). Either assay taken alone obscures this change
10 due to the wide excursion of hCG values which occur in normal pregnancy.

Others have documented progressive changes in hCG isoforms throughout pregnancy. Skarulis et al. found
15 that the fucose content of both intact hCG and also its free beta subunit increased as pregnancy progressed (Skarulis et al. 1992). Diaz-Cueto et al. investigating the isoelectric focusing pattern of circulating hCG throughout pregnancy, found that in early pregnancy,
20 more than 80% of the hCG isoforms were acidic. This fraction decreased to less than half (47%) late in the third trimester (Diaz-Cueto et al. 1996). In contrast, Wide and Hobson found that the hCG of early pregnancy was more "choriocarcinoma-like" by virtue of its greater
25 biological activity than the hCG of normal pregnancy (Wide and Hobson, 1987). Fein et al., in a study which employed gel filtration determined that first trimester hCG was a larger size than that of the third trimester. Treatment with exoglycosidases eliminated the size
30 differential, indicating that the first trimester hCG was more highly glycosylated (Fein et al. 1980).

001080"ST20E950

The B152/B109 ratio in matched serum/urine samples in the first and third trimesters of pregnancy compared with hCG from JAR cells.

5 The B152/B109 ratio in serum is analogous to that found
in matched urine samples and undergoes a similar change
as pregnancy progresses (Figure 5). The B152/B109 ratio
in the cell supernatant from JAR cells (a
choriocarcinoma derived cell line) was similar to that
10 of early pregnancy.

The B152/B109 ratios of both serum and urine hCG
concentrations are significantly higher in the first
15 trimester as compared to the third trimester of normal
pregnancies (Table 2). Significant differences between
serum and urine hCG concentration ratios as well as log
transformed ratios in early (5-6 weeks) and late (36-39
weeks) gestation were evaluated by paired t-tests (Table
3). In both the first and third trimesters, urinary
20 B152/B109 ratios were significantly higher than serum
ratios, indicating that there was a preferential
clearance of the B152-recognized isoform into urine,
regardless of the relative concentrations of the two
isoforms.

25

TABLE 2.

Analysis of the B152/B109 ratio in serum and in urine in the first vs third trimesters of pregnancy.

Measure	T-test (df)	P
Serum, ratio B152/B109	t(11)=6.65	0.0001
Serum, log(ratioB152/B109)	t(23)=21.61	0.0000
Urine, ratio B152/B109	t(11)=4.64	0.0007
Urine, log(ratioB152/B109)	t(15.7)=16.85	0.0001

007080" ST20E860

TABLE 3.

Analysis of the B152/B109 ratio in serum vs urine in the first and third trimesters of pregnancy.

Gestational age	Measure	Paired-t (df)	P
5-6 weeks	Ratio B152/B109	t(11)=3.25	0.0077
	Log(ratioB152/B109)	t(11)=6.25	0.0001
36-39 weeks	Ratio B152/B109	t(10)=5.47	0.0003
	Log(ratioB152/B109)	t(10)=7.14	0.0001

The B152/B109 ratio in urine samples from IVF patients

In urine samples from IVF patients (1-4 weeks post embryo transfer - ET) the B152/B109 ratio was again
 5 between 2-8 and decreased as pregnancy progressed (Figure 6), similar to that observed in natural conceptions. The effect of pregnancy duration with respect to outcome variables could best be represented by a linear or quadratic function. ANCOVA models including the second order week were fitted to the
 10 general equation: Outcome= (effect of time post ET)+(effect of diagnosis). After an appropriate ANCOVA model was determined, the least square means (adjusted for week post ET effect) were compared among the normal pregnancy, ectopic pregnancy and spontaneous abortion populations (Table 4). The log transformed values of
 15 both B109-B108* and B152-B207* measured hCG forms discriminated both ectopic pregnancy and spontaneous abortions from normal pregnancy (P=0.0001). The ratio of the log transformed values discriminated abortion from
 20 normal pregnancy (P=0.016). However, neither the ratio of B152/B109 nor the log of that ratio discriminated either of the pregnancy disorders from normal pregnancy.

007080" 5F20E950

A significant number of spontaneous abortions and ectopic pregnancies occur in IVF pregnancies. We did not find a difference in the ratio of the isoforms between either of these two categories as compared to normal controls, possibly a consequence of low statistical power. However a significant difference was found between the B152 hCG isoforms levels in normal pregnancy and spontaneous abortion. This supports our previous finding in early pregnancy loss, where diminished or absent levels of the B152 isoforms characterized an early pregnancy loss (O'Connor et al. 1998).

007080"5F20E960

TABLE 4.

IVF patients: analysis of covariance of hCG isoforms among normal pregnancy (np), ectopic pregnancy and spontaneous abortion as a function of gestational age.

Outcome	^c Adjusted R ²	^e -F	P	^d -Pairwise Difference
^a Log (ratio B152/B109)	0.51	0.89	0.41	none
^a Log (B109-B108*)	0.56	21.33	0.0001	np vs abortion & ectopic
^b Log (B152)/log (B109)	0.45	4.34	0.016	np vs abortion
^b Log (B152-B207*)	0.50	26.94	0.0001	np vs abortion & ectopic

- a- ANCOVA model with 2nd order polynomial coefficient (or parameter).
- b- ANCOVA model with only 1st order (linear) coefficient.
- c- Adjusted R² is a R² adjusting number of coefficients on the ANCOVA model so that comparisons of two R² with different ANOVA models with different number of coefficients are meaningful.
- d- "Pairwise difference" is based on t-test comparing the least-square means of outcome variables (after adjusting effect of week ET).
- e- Degree of freedom (df1, df2) for F-test are (2,82) for a model with only linear coefficient and (2,81) for a model with both linear and 2nd order coefficient.

HCG analysis of trophoblastic disease samples

5 Trophoblast disease serum (17 samples) and urine (28 samples) were obtained from patients post therapy and hence contained low hCG levels. Due to limited amounts of sample all of these specimens were run at a 1:10 initial dilutions. HCG levels in serum were low. The highest hCG concentration in serum was 202 fmol/ml in

001080"ST20E960

the B152-B207* assay, with a corresponding value of 148 fmol/ml in the B109-B108* determination. Six of seventeen samples in serum had detectable levels, with 4/6 having a higher value in the B152-B207* assay. Of the 15/28 positive urine samples however, 14/15 had higher levels in the B152-B207* assay than in the B109-B108* assay, with the highest hCG value being 20000 fmol/ml in the B152-B207* assay and 18715 fmol/ml in the corresponding B109-B108* assay. Due to the small sample size, no statistical treatment was performed on this data, but even in these post-treatment patients the B152/B109 ratio was ≥ 1 , which corresponds to the early pregnancy hCG isoform ratio.

The specimen limitations discussed above precludes our reaching any definitive conclusion on the analysis of trophoblastic disease samples. However it appears as might be anticipated that the B152 assay is more sensitive than B109 assay in detecting hCG immunoreactivity in the blood and in the urine of trophoblastic disease patients, even after treatment.

Chromatography of First Week of Gestation Pregnancy Pool. In order to determine whether the B152-B207* assay recognized other forms of hCG associated immunoreactivity in addition to the intact hCG molecule, specimens were pooled. FPLC on tandem Superose 12 columns followed by immunoassay of the fractions for all of the characterized forms of hCG revealed that only the intact hCG molecule (or hCG free beta subunit) gave a signal in this assay (See Figure 7). There were no lower molecular weight fragments identified by the B152-B207* assay. The hCG free beta analyte was measured in urine described in Figure 2 and was found to make a negligible contribution to over all hCG immunoreactivity in these specimens.

000000 "ST20E950

Molecules recognized by monoclonal antibody B152 in urine and pituitary extracts. In order to define the nature of the hCG isoforms recognized by B152, high resolution gel filtration columns of both pituitary extracts and postmenopausal urine concentrates were used. The rationale for use of pituitary extracts is to determine cross-reactive molecules, specifically those which are glycosylated, which are plentiful in pituitary which contains the entire family of glycoprotein hormones, hLH, hTSH, and hFSH as well as free subunits and the pituitary form of hCG. Two peaks are detected in both of these cases. Only one peak was detected in similar studies of pregnancy urine concentrates as described earlier. In the pituitary, it is likely that the larger molecule is pituitary hCG (70K) while the smaller sized molecule is hLH. Since hLH is present at 100x or so as compared to pituitary hCG, the apparent similar concentration of immunoreactivity indicates that B152 has reduced cross-reactivity to hLH as compared to hCG. Likewise, both hCG and hLH occur in postmenopausal urine, again with much more hLH than hCG and the B152 pattern is similar to that of the pituitary extract. These results show that B152 is generally hCG specific except for cross reactivity to hLH (as shown by standard cross-reaction studies in Table I) and that its carbohydrate specificity is both to the protein portion as well as to the carbohydrate moieties of hCG (and to a lesser extent of hLH) since it does not react with the multitude of other glycoylated proteins present in the pituitary nor with those in postmenopausal urine except for hCG or hLH-related molecules.

Serum and urine specimens were analyzed using two assays, B109-B108* and B152-B207*, which recognize the difference in molecular isoforms of hCG. See Table I. The *in vitro* bioassay for hLH/hCG is described above. (See Figure 1). The immunometric assay employs 96-well

998038

microtiter plate technology. The coating antibody, at a concentration determined to provide the most satisfactory combination of sensitivity and range, is applied to the microtiter wells (Immulon IV, Dynatech Laboratories) in carbonate buffer (0.2M, pH 9.5). The plates are incubated with the coating solution at 4°C, overnight, then aspirated, washed with washing solution (0.05% Tween, 0.15N NaCl), and blocked with a 1% solution of BSA (three hours at room temperature). The BSA solution is aspirated and the appropriate hCG standards (200µL/well), in buffer B (PBS/0.1% bovine IgG/0.1% sodium azide), or in hCG free serum (Chemicon, Inc.), or hCG free urine, as appropriate to the specimen matrix, and specimens are added to the wells. The plates are sealed with plate sealers, and incubated overnight at 4°C. The controls, specimens, and standards are then aspirated, the plates washed 5 times with washing solution, and iodinated detection antibody in buffer B (200 uL/well, 100,000 cpm/well) added and incubated overnight at 4°C. The wells are again aspirated, washed 5 times with washing solution, separated and counted (Packard Cobra gamma counted). Values are interpolated from a smoothed spline transformation of the count data. This assay procedure, as well as assay validation has been previously reported (O'Connor, J. F., et al., 1988).

Creatinine analysis, when urine values are normalized to creatinine, is performed in a microtiter plate format following a modification of the Tausky procedure (Tausky, H. H., 1954).

Descriptive statistical and graphical methods are applied to measures of serum and urine samples from normal healthy pregnancies to identify the distributions a) between patient first trimester average B152 levels, B109 levels and B152/B109 ratio; b) between patient

007080-5720E960

variability in time to B152/B109 ratio reaching 1.00; and c) between patient variability in time to B152/B109 ratio declining by 1/3rd from first trimester maximum levels. The variability in the timing of the crossover
5 in the ratio of these two analytes provides an empirical basis from which to estimate the value of these markers as biochemical signatures of a viable third trimester fetus.

10 Comparison of the assay profile of healthy normal pregnancies to those of unsuccessful pregnancies from failed IVF implantations, two non-parametric hypotheses are available: 1) the proportion of pregnancies in which the B152/B109 ratio falls below 1.00 is no different in
15 healthy normal and unsuccessful IVF pregnancies; 2) the proportion of pregnancies in which the B152/B109 ratio declines by 1/3rd from first trimester maximum levels is no different in healthy normal and unsuccessful IVF pregnancies. These hypotheses can be tested as a
20 difference between two proportions. For example, a comparison of week 14 vs. week 9, week 13 vs. week 6, week 12 vs. week 5 or week 11 vs. week 4 pregnancies to show a reversal of the B152/B109 ratio in healthy normal pregnancies and unsuccessful IVF implantations,
25 respectively. The power analyses apply to an outcome defined as the time at which the B152/B109 ratio declines by 1/3rd from first trimester maximum levels, although this outcome would necessarily provide earlier detection of pregnancy failure than the reversal of the
30 B152/B109 ratio. Patterns of results less discriminantly different from these indicate a rejection of the dichotomous outcome of B152/B109 ratio reversal as a clinically meaningful marker of pregnancy failure.

35 Alternatively, the same two non-parametric hypotheses can be recast as parametric hypotheses by considering the timing of the biochemical events within the assay

profile of healthy normal pregnancies and unsuccessful pregnancies from failed IVF implantations: 1) the time at which the B152/B109 ratio falls below 1.00 is no different in healthy normal and unsuccessful IVF pregnancies; 2) the time at which the B152/B109 ratio declines by 1/3rd from first trimester maximum levels is no different in healthy normal and unsuccessful IVF pregnancies. Of course, the objective is to provide an empirical basis from which clinicians may counsel their patients. Thus, it is important to adopt a logistic model for this component of the data analysis. With pregnancy success as the outcome, logistic models allow the estimation of the (symmetrical) hypothesis of increase in risk of pregnancy failure for each additional week where either the B152/B109 ratio has failed to decline by one third from first trimester baseline maximum values or the B152/B109 ratio has failed to become less than 1.00 (measured in weeks). The logistic model enables specification of the time at which results indicate a particular pregnancy exceeds an a priori defined likelihood of failure, given assay data regularly available during pregnancy, and allows incorporation of other risks for pregnancy failure in the same data analytic framework to assess the relative contribution of threats to pregnancy loss. The Cox proportional hazard model may be used to examine predictors of the crossover rates. Mixed effects models can also analyze repeated measures of the B152/B109 ratios taken during entire cycles. These models are particularly useful since they allow inclusion of incomplete and imbalance data (i.e. data with missing values and unequal timing of data collection), to estimate effects of time-varying covariates, to model dependency structure of repeated measures and to model possible heterogeneity of the ratio measures within each experimental group.

00T080"ST20E960

B152 hCG isoforms isolated from early pregnancy urine and determination of their protein and carbohydrate structures.

5 hCG molecules are isolated from urine collected from women in early pregnancy for both protein and carbohydrate analyses. According to one approach, molecules are isolated from HPLC fractions, digested with proteases before and after reduction of disulfide
10 bonds, examination of the resultant peptides by mass spectrometry and/or sequence analysis, isolation of carbohydrate moieties after glycosidase digestions and determination of carbohydrate structures by a combination of specific glycosidases and retention times
15 on specialized anion exchange columns as compared to know branch-chain oligosaccharide standards. In a similar approach, the final purification stage for the isolated hCG isoforms is SDS gel electrophoresis. Both protease digests and glycosidase digests are performed
20 on the blotted and cutout band. This method results in greater purity of the protein and less artifactual errors due to contamination by carbohydrates which are not in the purified protein but are derived from outside contaminants.

25

Carbohydrate compositional analyses and oligosaccharide branched chain identifications.

The MALDI TOF mass spectrometric method may be used to confirm oligosaccharide structures by using specific
30 glycosidases on the glycopeptides and determining the change in molecular weight as the sugars are digested off the glycopeptide. Only the hCG beta COOH peptide can be expected to contain O-linked sugar moieties. These are of special interest since it is thought that
35 B152 has significant reaction with this region. The structures of this region can be determined in a similar fashion using enzymes that specifically release O-linked

000000" ST 200000

glycans. The O-linked structures has been previously examined using standard reference pregnancy hCG (Cole, L. A., et al., 1985). The O-linked branched chain structure are determined by a similar strategy using the
5 Dionex chromatographic system as well as specific glycosidases on the C-terminal glycopeptides and Mass Spectrometry. In one study (Elliott, M. M., et al., 1997), these techniques were used to elucidate the carbohydrate structures of CR series hCG preparations
10 (standard urinary pregnancy hCG) and compared them to the structures of patient samples such as C5 which was the immunogen employed to generate antibody B152. It was found that C5 contained significantly more mono and tri-antennary (2X mono and 3X tri-structures than the CR preparations)
15 (on the N-Asn residues. It was also found that more tetrasaccharide structures were on the hCG COOH-terminal peptide O-Serine residues in the choriocarcinoma hCG isoform than in the CR preparations.

20 Biological activity and metabolic clearance of hCG isoforms. Biological activity is a function both of molecular structure and half-life in the circulation, which can be influenced by structure. Alterations in
25 carbohydrate/sialic acid content of the glycoprotein hormones are thought to be responsible for the changes in hCG biological/immunological activity observed throughout pregnancy. In addition, signal transduction at the receptor is influenced by the pI of the hCG
30 isoform and the presence or absence of carbohydrate. Thus, it is valuable to examine both receptor binding and biological activity *in vitro* and, in order to determine the mechanism of action, to distinguish receptor binding and signal transduction as well as
35 relative potency of signal transduction along with *in vivo* bioactivity determinants such as circulating half life. Studies, including clearance rates, are performed

001000"ST200960

on B152 hCG isoforms of early successful pregnancy, hCG from third trimester pregnancy, and the reference urinary hCG preparation, CR 127.

5 Example 3: B152 and B151 immunoreactivity in non-trophoblastic malignancy.

With the exception of trophoblastic disease and testicular cancer, hCG is expressed in the blood of about 20% of patients with all other types of cancer (Hussa, R. O., 1987). HCG beta core fragment in the urine has a significantly higher level of expression, especially in gynecological malignancy. Since the B152 antibody was developed to a form of hCG produced in a malignancy, it was of interest to examine the expression of B152 and nicked hCG immunoreactivity (B151) in non-trophoblastic malignancy. Accordingly, blood and urine derived from men undergoing chemotherapy for prostate cancer or women for ovarian cancer were evaluated for the expression of hCG isoforms in plasma and urine. It is significant that in prostate cancer, B152 hCG immunoreactivity is found in the blood and urine of prostate cancer patients in instances when there is no hCG detected by B109-B108*. In ovarian cancer patients evaluated, there is evidence of nicked hCG in the blood, even in the absence of both B109 and B152 immunoreactivity. Neither of the above groups demonstrated the presence of hCG immunoreactivity when the standard pregnancy derived hCG assay was employed. It is reassuring to find that nicked hCG, the existence of which has been documented by several investigators, can be found and reliably measured in a clinical setting.

35 Experimental Discussion

In the course of these studies, a potentially important new signal was observed in the urine of women early in pregnancy, namely an epitope of a form of hCG which may

007080" ST20E960

indicate the likely success of carrying a pregnancy. Likewise, absence of this signal may indicate that EPL will occur. Since EPL can be a very sensitive marker of environmental toxins (Hakim, R. B., et al., 1995) and is frequently used as an epidemiological marker of exposure, the finding of this epitope provides a powerful tool for monitoring the safety of the environment. In addition, this assay facilitates increasing the success rate of IVF infertility programs since the predictive value of the new measuring system would rapidly indicate successful approaches. Described herein is the novel and completely unexpected finding that successful pregnancies display a high content of unique isoforms of hCG that are maintained for the first few weeks of pregnancy and then rapidly decline as pregnancy progresses. Based on properties of the immunoassay system, it is hypothesized that these hCG isoforms may be hyperglycosylated. This is a striking observation never reported nor suspected earlier. Carbohydrate analyses (Elliot, M., 1997) demonstrate that C5 hCG employed as immunogen for antibody B152, contains two times the monoantennary content and three times the tri-antennary content of branch chain sugars as compared to the CR series of natural pregnancy urinary hCG. In addition, the O-linked carbohydrates are mostly tetrasaccharide instead of disaccharide in C5 as compared to CR 127 hCG. (CR 127 hCG is similar to the WHO preparation, the third international hCG standard, which was CR 119 hCG, prepared by Canfield and Birken twenty years ago but still in use today) (Birken, S., et al., 1991a). B152 recognizes C5 hCG much better than nicked CR127 hCG or non-nicked CR 127 hCG (Birken, S., et al., 1993). In addition, JAR cell type hCG is known to contain a similar array of carbohydrate moieties. It was found to be recognized by B152 similar to the early pregnancy isoforms in healthy pregnancies. The observation that the hCG isoform produced by JAR cells

001080"ST20E960

in culture (B152/B109 ratio) is similar to that found in early pregnancy hCG isoforms supports the hypothesis that the production of a type of hCG with a particular glycosylation pattern is a prerequisite for a viable pregnancy. This glycosylation pattern is not characteristic of the hCG of later pregnancy.

A variety of pregnancy disorders are testable. One category of patients consists of those women who experience a high rate of recurrent abortions. Even in populations with no known fertility problems, the total rate of pregnancy loss is 32% (EPL plus clinically recognized abortion) (Wilcox, A. J., et al., 1988). The risk of recurrent abortion increases with the number of spontaneous abortions experienced in the past, reaching an incidence of 32% after three consecutive abortions. (Hill, J. A., and Anderson, D. J., 1990). Probable causes of recurrent spontaneous abortion, comprising genetic, infectious, hormonal imbalance, or immunologic factors can be established in less than 60% of all spontaneous abortions, leaving 40+% of spontaneous abortions with a completely unestablished etiology. These facts, taken together with reports establishing that the administration of exogenous hCG can be an effective therapy in subjects with a history of recurrent spontaneous abortion (Quenby, S., and Farquharson, R. G., 1994; Harrison, R. F., 1985) lends support to the hypothesis that a disproportionate production of the ineffective isoforms of hCG in early pregnancy is a causal factor in both early pre-clinical loss as well as in spontaneous abortion.

A second category includes women undergoing embryo transfer. These patients provide several distinct advantages: The patients undergoing this procedure are not treated with crude hCG preparations, making measurement of hCG isoforms easy and decisive since all

001080"ST209660

hCG forms derive from the embryo none from any injected hCG preparations. Second, is the opportunity to monitor the nature of the isoforms from day 9 of a successful pregnancy. Third, is the ability to obtain large volumes of urine to purify the early pregnancy isoforms to determine their structures. Fourth, since pregnancy loss is from 50% to 70% in this population, the loss can be defined as due to lack of the essential hCG isoform recognized by B152 or due to other causes. Comparison of early pregnancies in populations of women not undergoing *in vitro* fertilization procedures with those undergoing embryo implantation can, thus, assess whether pregnancy loss situations present similar or different patterns of hCG isoforms during the process. The mechanism of pregnancy loss in the general population as compared with the much higher rate of embryo loss in IVF programs may be different. Additionally, it has been established that the hCG produced in choriocarcinoma has differences in carbohydrate structures, sialic acid content and biological activity (Wide, L., and Hobson, B., 1987; Elliot, M., et al., 1997; Husa, R. A., 1987). Since B152-B207* assay incorporate monoclonal antibodies raised against an immunogen derived from choriocarcinoma, specimens may be evaluated from patients with gestational trophoblastic disease in order to determine whether the above assays recognize the hCG produced in these conditions with greater sensitivity and specificity than do assays based on the hCG of normal pregnancy, as is apparently the case for the hCG produced in testicular and ovarian cancer.

There are few reports of changes of carbohydrate content of hCG-related molecules during pregnancy. Blithe and colleagues studied free alpha subunit of hCG whose carbohydrate content differs from that of alpha within hCG by additional carbohydrate antennae and fucose. The carbohydrate of free alpha becomes increasingly complex

09630215 "030100

in terms of more branches and higher content as pregnancy proceeds. It has also been reported that the quantity of fucose increased in both hCG and in free alpha as pregnancy proceeded (Skarulis, M. C., et al., 1992). Thus, the literature indicates increasing content and complexity of carbohydrate of hCG and free alpha subunits. However, immunological data using the B152 monoclonal antibody, implies a progression to simpler carbohydrate content during pregnancy. Since the beta COOH-region's O-linked carbohydrates may be involved in the epitope recognized by B152, it is conceivable that the carbohydrate structures of this region may be altered in a different pattern from the N-linked glycans studied by Blithe and colleagues (Skarulis, M. C., et al., 1992; Blithe, D. L., and Iles, R. K., 1995). Data from Skarulis et al. indicate that heterodimeric hCG may contain additional fucose but do not provide data that this late pregnancy hCG becomes hyperglycosylated as does free alpha.

Other studies indicated that the forms of hCG during EPL likely differ in biological activity from those hCG isoforms in successful pregnancies (Ho, H.-H., et al., 1997). The *in vitro* bioassays employed in those studies are unsuitable for large-scale studies and are not as reliable as the immunoassays described herein. Furthermore, it is likely that *in vivo* assays may give different results since *in vitro* and *in vivo* assays sometimes give completely disparate results. In this case, *in vivo* and clearance assays are most important in order to identify whether the hCG isoforms are truly more potent in the whole animal and to identify the reasons for the increased potency. Thus *in vitro* and *in vivo* bioactivities of the early pregnancy isoforms of hCG are highly significant.

Carbohydrate differences is a widely accepted

001030"ST202960

5 explanation for variations in biological to
immunological ratio such as the forms observed by
various studies of EPL (Ho, H.-H., et al., 1997).
10 Various studies (Grotjan, H. R. J., and Cole, L. A.,
1989; Hoermann, R., 1997; Stanton, P. G., et al., 1993;
Szkudlinski, M. W., et al., 1995, Thotakura, N. R., et
al., 1994; Szkudlinski, M. W., et al., 1993), have shown
that sialic acid differences are an explanation for such
heterogeneity in biological activities of glycoprotein
15 hormones. These studies have also confirmed the dogma
that *in vitro* biological activities can yield the
opposite results from *in vivo* studies because of altered
metabolic clearance rates in the latter studies. Thus,
more acidic (more highly sialylated) forms of
20 gonadotropins are more biopotent in the whole animal
because of prolonged circulating half-lives. The same
molecules may appear less potent in *in vitro* assays due
to greater acidity, greater negative sialic acid
content. Hoermann et al. (Hoermann, R., et al., 1997)
25 demonstrated the exclusion of many of the acidic
circulating hormone forms from the urine, thus,
prolonging their half-lives. The pI pattern of normal
pregnancy as well as trophoblastic cancer hCG in serum
is quite different from that of urine. Since the
30 studies described herein indicate that EPL hCG isoforms
have reduced *in vitro* biological activity, this finding
cannot be explained solely by what is known of
biological activity and sialic acid content. Early
pregnancy isoforms recognized by monoclonal antibody
35 B152 may be more potent *in vivo* by virtue of prolonged
half-life they may then display increased signal
transduction at the receptor as well. This may be
explained by a hyperglycosylated form of hCG which is
not hypersialylated. In this case, the extra sugar
portion would help prolong circulating half-life of a
more basic pI form of hCG which also has increased *in
vitro* bioactivity.

00000000000000000000000000000000

Example 4: Diagnosis of Gestational Trophoblast Disease.

5 An important application of the B152 (early hCG isoform)/B109 (late hCG isoform) ratio analysis described herein above is in the very early (and facile) diagnosis of gestational trophoblast disease. Examples of gestational trophoblast disease include choriocarcinoma or hydatidiform mole. In normal pregnancy, the ratio of B152/B109 of the two isoforms of hCG rapidly decreases, eventually inverting. In gestational trophoblast disease including choriocarcinoma or hydatidiform mole, the ratio is initially higher than found in normal pregnancy, but does not diminish during the course of the apparent pregnancy. This approach provides a highly sensitive and specific diagnostic marker for gestational trophoblast disease.

20 Other pregnancy disorders in which hCG levels are abnormally high or abnormally low include Down's syndrome or other aneuploid pregnancies, ectopic pregnancy, preeclampsia, and intra-uterine growth retardation. Because the hCG production in these conditions is quantitatively abnormal compared with normal pregnancy, an altered ratio of the hCG isoforms identified by B152 (early hCG isoform) and B109 (late hCG isoform) can be detected.

30 Thus, the dual isoform analysis (B152/B109) further provides a method for diagnosing pregnancy disorders and gestational trophoblast disease.

Materials and Methods

35

Hormones

The non-nicked hCG isolated from the CR127 preparation

09530215-00000

of hCG was used as a standard in both assays (Birken et al. 1993). The pituitary hCG was isolated as described (Birken et al. 1996). C5, a 100% nicked hCG having extra sugars on both N- and O-linked carbohydrate moieties, 5 purified from the urine of a choriocarcinoma patient (Elliott et al. 1997), was supplied by Dr. Laurence Cole (Yale University School of Medicine). Although the C5 immunogen used in the development of B152 antibody was 100% nicked hCG isoform (i.e. had cleavages in the 10 peptide backbone of loop 2 of the b subunit) the antibody did not discriminate nicked from non-nicked hCG (O'Connor et al. 1998).

The same serial dilutions of non-nicked hCG, pituitary 15 hCG and C5 were used for binding characterization in hCG assays. Hormone concentrations of initial stock standards solutions were determined by amino acid analysis.

20 Immunoradiometric assays (IRMA)

The methodology used in the construction and validation of the ¹B109-B108* assay has been fully described elsewhere (O'Connor et al. 1988). The B152-B207* assay 25 has also been characterized (O'Connor et al. 1998). Both assays were performed with a slight modification of the published procedure: the capture antibody was adsorbed onto the wells of microtiter plates (Immulon IV, Dynatech, Chantilly, VA) by incubating a 5 µg/ml 30 solution (B109-B108* assay) or 25 mg/ml solution (B152-B207* assay) in coating buffer (0.2 M bicarbonate, pH 9.5) overnight at 4 C. The coating antibody solution was aspirated, the plates washed (wash solution: 0.9% NaCl, 0.05% Tween 20) and blocked with a 1% solution of BSA in

35 The first antibody is a capture, the second antibody with an asterisk is an iodinated detection antibody.

098030 "STRE960

PBS with 0.1% sodium azide. Following incubation with the BSA solution (minimum 3 hours at room temperature) the blocking solution was removed, the wells again washed with wash solution and 200 ml/well of the appropriate hCG standards were added in phosphate buffer B (PBS with 0.1% bovine gamma globulin and 0.1% sodium azide). After overnight incubation at 4 C, the plates were again aspirated and washed. The 200ml (50,000 cpm-100,000 cpm) of ¹²⁵I- labeled antibody was added to the wells which were again incubated for 24 h at 4 C. The tracer was aspirated, the plates washed with wash solution, the individual wells placed in glass tubes and the radioactivity determined in a Packard Cobra gamma counter. Doses were determined by interpolation from a smoothed spline transformation of the data points.

All samples were stored frozen at -20 C prior to assay. Because extreme values of sample pH may interfere with antibody binding, the urine pH was adjusted with 1.0M Tris (pH 9.0), 50µl/ml urine prior to assay, so that the final pH was in the range of 7-7.4 (O'Connor et al. 1988). Intra-assay variation was 6% for both assays, inter-assay variation was 12% for B109-B108* and 13% for B152-B207* assays. Sensitivity (least detectable dose) defined as +2SD from the zero calibrator, was 1 fmol/ml for the B109-B108* assay and 2.2 fmol/ml for B152-B207* assay.

Patients samples

Urine samples from IVF patients were a gift from Dr. L. Cole. They included spontaneous abortion (n=14, range of gestational age 1.8-4.1 weeks from ET- embryo transfer), ectopic pregnancies (n=7, gestational age 2.3-4 weeks) and normal pregnancy controls (n=65, encompassing the range 0.6 to 5.4 week from ET). Some of the normal pregnancy urine samples throughout the pregnancies were

001080 " STREETS 0

also obtained from Dr. Cole. Others were obtained from the clinical practice of collaborating physicians at Columbia Presbyterian Medical Center (CPMC) (Total n=103). Matched serum/urine samples from the first 5 (n=12) and the third (n=11) trimesters were provided by Dr. Amalia Kelly at CPMC. Trophoblast disease serum (n=17) and urine (n=28) samples were obtained from Dr. Cole, but were collected by Dr. Edward Newlands (Charing Cross Hospital, London, UK). All specimen 10 collection protocols were approved by the appropriate Institutional Review Board.

Statistical Analysis

15 Polynomial regression models of log transformed hormone ratios were used to describe the relationship between the change in ratio as a function of gestational age in normal pregnancy. A paired t-test was used to evaluate the relationship between matched serum and urine hormone 20 ratios. Analysis of covariance (ANCOVA) was used to describe the time adjusted relationship of hormone values in ectopic pregnancy and spontaneous abortion to those of normal pregnancy.

Urine Processing.

25 Twenty-four hour urine samples are collected from women undergoing embryo transfer as well as women in early natural pregnancy. The urine is refrigerated during the collection procedure. After delivery of the urine to 30 the laboratory, sodium azide is added to 1g/liter. Women undergoing *in vitro* embryo transfer are not pre-treated with hCG. Thus, all hCG which appears in their blood or urine is derived from the embryo (except for the small amounts of pituitary hCG present in all 35 people). Raw urine is freed from particles by centrifugation followed by Pellicon filtration through a 0.45 micron membrane. Next, the procedure is to

001080" 5120E960

concentrate the urine with a Pellicon (Millipore) system which concentrates as much as 30 liters to 500ml overnight (4°C) using a 3,000 MW cutoff membrane. Smaller volumes can be concentrated in less than two hours. Next, the urine is desalted and delipidated by passage through a large volume of Sephadex G25 in 0.1 M ammonium bicarbonate. This step greatly increases the binding of CG to immunoaffinity columns. The desalted urinary concentrate is next size fractionated on the Pharmacia HiLoad Superdex 200 and the hCG and hCG subunit peaks are identified by specific immunoassays (O'Connor, J. F., et al., 1994) and the appropriate fractions are pooled and dried. The hCG and hCG subunits are purified from the gel filtered urine concentrate by immunoaffinity on insolubilized hCG antibody columns as described but with the use of either 4M guanadine (0.1M tris acetate, pH 5) or ammonium thiocyanate as eluant to decrease loss of sialic acid from the hormone. Alternatively, hCG is purified by conventional chromatographic procedures, anion exchange and hydrophobic chromatography. The subunits are separated on reverse phase HPLC using a 0.01M sodium phosphate, pH 5 buffer and acetonitrile, after incubation in 4M guanadine, 0.1M tris acetate, pH 5. A third method is final purification and separation of the hCG subunits on SDS PAGE electrophoresis followed by electroblotting to PVDF. The PVDF band can be subjected to protease digestion to release peptides and glycopeptides which can be separated on reverse phase HPLC in neutral pH 5 buffers.

Separation of Glycopeptides from Isolated hCG subunits.

To facilitate isolation of the glycopeptides from the hCG subunits, the subunits are both tryptic digested and the products of digestion are separated on reverse phase HPLC (using a pH 5 buffer). This procedure results in removal of the large beta COOH-terminal peptide which

0950215 080100

contains O-linked sugars. It also releases small, non glycopeptides from both subunits (Pollak, S., et al., 1990, Birken, S., et al., 1987; Birken, S., et al., 1986). Next, the main disulfide-linked core of each hCG subunit, is reduced and carboxymethylated, and separated on reverse phase HPLC at pH 5. At this stage, large peptides are isolated, including the glycopeptides. Each separated glycopeptide is redigested with trypsin and re-separated on HPLC at pH 5. These glycopeptides are next employed for two different methods of sugar chain analysis. One method is the approach of releasing the oligosaccharides by enzymatic digestions using PNGase F for the N-linked glycans. The released glycans can be separated from the peptides by ethanol precipitation, desialyated with neuraminadase, and separated directly on a Dionex Carbopac PA-100 column. Oligosaccharide standards are available from Dionex, Oxford Glycosystems and other companies for calibrating column elution times for various glycans (Hardy, M. R., and Townsend, R. R., 1994, Rohrer, J. S., et al., 1993, Weitzhandler, M., et al., 1993; Townsend, R. R., et al., 1989). Confirmation of the released structures is obtained by performing carbohydrate compositional analysis of eluted glycan peaks as well as performing digestions with specific glycosidases and rechromatographing the modified glycan on the Dionex system (Hardy, M. R., and Townsend, R. R., 1994; Rohrer, J. S., et al 1993; Weitzhandler, M., et al., 1993; Townsend, R. R., et al., 1989; Townsend, R. R., et al., 1991; Townsend, R. R., et al., 1989; Hardy, M. R., and Townsend, R. R., 1989; Townsend, R. R., et al., 1988; Hardy, M. R., et al, 1997; Hardy, M. R., and Townsend, R. R., 1988; Dionex, 1997; Spellman, M. W., 1990; Kumarasamy, R., 1990). The newly modified glycan can be observed to elute at the same time as the appropriate standard oligosaccharide and, in addition, the released monosaccharide can frequently be identified as well (Dionex, 1997). Structure determination is

095901510000

facilitated by the use of specific glycosidases for branch chain cleavage as well as for digestion of individual sugars from each of the branch chains. For example, Endo H cleaves high mannose type and hybrid oligosaccharide chains while glycosidase Endo F2 cleaves biantennary complex types and PNAase F cleaves tri and tetra-antennary chains down to the N-Asn bond.

Competitive receptor binding and in vitro bioassay.

Bioassays are performed with recombinant-engineered CHO cells transfected with the human receptor to LH/CG Cells are maintained in Ham's F-12 medium, 4mM Glutamine, 400ug/ml G418 (Gibco), 5% fetal calf serum, 100IU/ml penicillin, 100ug/ml streptomycin. The cells are removed from the flask surface by versene only.

A competitive receptor assay constructed as follows: The receptor binding assay mixture contains 100 ul of the appropriate dilution of serum/urine samples or hCG dilutions for standard curve, 100ul of ¹²⁵I-hCG (50,000-100,000cpm) in buffer A(PBS/0.1%BSA) and 100ul of CHO cells (2x10⁵ cells in PBS). The mixture is incubated at 37°C with slight shaking followed by centrifugation for 10 minutes at 750xg. The supernatant is aspirated and the cell pellet is counted in gamma-counter.

In vitro bioassay. Transfected CHO cells are seeded (200,000 cells/well) into a 24 well plate in culture medium and incubated for 2-3 days until the cells reach confluence. Non-transfected CHO cells are included to monitor non-specific response. The medium is removed and replaced with medium containing 1 mM isobutylmethylxanthine with appropriate dilutions of tested serum or urine. The plates are incubated at 37°C for two hours. The supernatant is removed, and the wells washed with Hank's balanced salt solution. The intracellular cAMP is extracted with 95% ethanol, which

09630215 "090100

is diluted 1:5, (or up to 1:40, depending on cAMP content) in assay buffer provided by the cAMP kit (New England Nuclear). cAMP assay is performed according to manufacturer's instructions. Response is normalized to well protein content (BCA protein assay kit, Pierce, Rockford, IL).

In vivo bioassay is determined by the uterine weight assay in immature female mice, following the procedure of Wide and Hobson (Wide, L., and Hobson, B., 1987). The mice are injected subcutaneously with one third of the total dose of gonadotropin on three consecutive days and killed 72 hours after the first injection. Uteri are dissected free from mesentery, fat and oviducts, blotted to remove intrauterine fluid and weighed to the nearest 0.1mg. Five to ten mice are used at each of these dose levels. The hCG standard preparation used is a nicked hCG. This material may be run concurrently with specimens isolated from first and third trimester pregnancy. Sham saline injection may be used as a control. The response signal is the log mouse uterine weight.

Clearance of hCG isoforms. The clearance of hCG is determined in the rat. Blood (200ul/sample) is obtained at 0, 120, 240, 360 and 480 minutes post injection, from an indwelling catheter in an catheterized external jugular vein, following the procedure described by Newman et al. (Newman, C. B., et al., 1985) and Brown and Hedge (Brown, M. R., and Hedge, G. A., 1972). Briefly, adult male Sprague-Dawley rats (Charles River Laboratories, Wilmington MA), wt 175-225g, are given free access to food and water. Rats are handled for acclimatization for one week after arrival, and several days before the hCG infusion, the rats are cannulated under pentobarbital anesthesia. A 21 gauge stainless steel cannula is inserted into the one external jugular

0953025 090100
001000 "5 F20E960

vein. The placement of the catheter allows for the collection of blood from the unrestrained, unstressed rat. After the animals have recuperated from the cannula implacment , an hCG isoform is injected (10 5 μ g/ml sterile saline) through the cannulated vein. Blood samples are obtained at the four time intervals listed above. The blood is allowed to clot and the serum separated and stored at -80°C for immunometric assays specific for different hCG isoforms.

10 Clearance rate of the isoforms of hCG from the circulation of the rat are estimated by computer fitting the concentration data to an equation of the general form:

15 Concentration= $Ae^{-\alpha t} + Be^{-\beta t}$ at time t ; A and α are parameters of the rapid component and B and β are parameters of the slow component. The metabolic clearance rate (MCR) is calculated as $\text{MCR} = \text{Dose}/(A/\alpha + B/\beta)$ and the initial volume of distribution is 20 calculated from $V_d = \text{Dose}/(A+B)$. The MCR is normalized to body weight for statistical analysis, which is performed using ANOVA with Duncan's range test for determination of significance (Cassals, J. W., et al., 1989).

25 Mice. The mouse species used in the experiments described herein are Balb/c mice, aged 12-20 weeks old and adult Sprague-Dawley rats of either sex. Mice used for the production of monoclonal antibodies through ascites and for the determination of *in vivo* biological activity as described. Balbc/c mice are used because 30 hybridoma cell lines were developed using Balb/c splenocytes.

000000 "5000000000

References

- Amano, J., R. Nishimura, S. Sato, and A. Kobata. 1990. *Glycobiology*. 1:45-50.
- 5 Armstrong, E. G., P. H. Ehrlich, S. Birken, J. P. Schlatterer, E. Siris, W. C. Hembree, and R. E. Canfield. 1984. *J. Clin. Endocrinol. Metab.* 59:867-874.
- Baenziger, J. U. 1994. *FASEB J.* 8:1019-1025.
- 10 Bahl, O. P., L. Marz, and W. R. Moyle. 1995. pp. 125-44. In Anonymous In: Dufau ML, Means AR, ed. Hormone binding and target cell activation in the testis. New York, Plenum Press, 1974.
- 15 Berger, P., Schwarz, S., Spottl, G., Wick, G. and Mann, K. (1993) Variants of human chorionic gonadotropin from pregnant women and tumor patients recognized by monoclonal antibodies. *J. Clin. Endocrinol. Metab.* 77, 20 347-351.
- Birken, S., M. A. Kolks, S. Amr, B. Nisula, and D. Puett. 1987. *Endocrinology* 121:657-666.
- 25 Birken, S., Gawinowicz, MA, Kardana, A., Cole, LA. 1991. *Endocrinology* 129: 1551-1558.
- Birken, S. , Kovalevskaya, G. , O'Connor, J. *Mol Cell Endocrinol*, 1996, 125:121-131.
- 30 Birken, S., M. A. Gawinowicz Kolks, S. Amr, B. Nisula, and D. Puett. 1986. *J. Biol. Chem.* 261:10719-10727.
- 35 Birken, S., Y. Chen, M. A. Gawinowicz, J. W. Lustbader, S. Pollak, G. Agosto, R. Buck, and J. O'Connor. 1993. *Endocrinology* 133:1390-1397.

09530215 030100
007080 "ST0E960

- Birken, S., Y. Maydelman, M. A. Gawinowicz, A. Pound, Y. Liu, and A. S. Hartree. 1996b. *Endocrinology* 137:1402-1411.
- 5 Blithe, D. L. and R. K. Iles. 1995. *Endocrinology* 136:903-910.
- Braun, J. R., T. E. Willnow, S. Ishibashi, G. Ashwell, and J. Herz. 1996. *J. Biol. Chem.* 271:21160-21166.
- 10 Brown, M. R. and G. A. Hedge. 1972. *Neuroendocrinology*. 9:158-174.
- Browne, E. S., M. V. Flasch, M. R. Sairam, and V. K. Bhalla. 1990. *Biochim. Biophys. Acta* 1033:226-234.
- 15 Cassals, J.W., Mann, K., Blithe, D.L., Nisula, B.C., Wehmann, R.E. 1989. *Cancer* 64:2313-2318.
- Chang, P.L., Canfield, R.E., Ditkoff, E.C., O'Connor, J.F., Sauer, M.V., 1998. *Fertil.Steril.*, 69:412-414.
- Chmielewski, C. 1992. *ANNA. J.* 19:34-38.
- 25 Cole, L.A. and Kardana, A. (1992) Discordant results in human chorionic gonadotropin assays. *Clin. Chem.* 38, 263-270.
- Cole, L. A., A. Kardana, P. Andrade-Gordon, M. A. Gawinowicz, J. C. Morris, E. R. Bergert, J. O'Connor, and S. Birken. 1991a. *Endocrinology* 129:1559-1567.
- 30 Cole, L. A., A. Kardana, F. C. Ying, and S. Birken. 1991b. *Yale J. Biol. Med.* 64:627-637.
- 35 Cole, L. A., A. Kardana, S. Y. Park, and G. D. Braunstein. 1993. *J. Clin. Endocrinol. Metab.*

76:704-710.

Cole, L. A., S. Birken, and F. Perini. 1985. *Biochem. Biophys. Res. Commun.* 126:333-339.

5

Diaz-Cueto, L., Barrios-de-Tomasi, J., Timossi, C., Mendez, J.P. and Ulloa-Aguirre, A. (1996) More in-vitro bioactive, shorter-lived human chorionic gonadotrophin charge isoforms increase at the end of the first and during the third trimesters of gestation. *Mol. Hum. Reprod.* 2, 643-650.

10

Dionex. 1997. *Technical Note* 42

15

Elliott, M. M., A. Kardana, J. W. Lustbader, and L. A. Cole. 1997. *Endocrine*, 7:15-32.

Ellish, N.J., Saboda, K, O'Connor, J, Nasca, P.C., Stanek, EF, Boyle, C. *Hum Reprod* 1996 11:406-412.

20

Fares, F. A., N. Suganuma, K. Nishimori, P. S. LaPolt, A. J. Hsueh, and I. Boime. 1992. *Proc. Natl. Acad. Sci. U. S. A.* 89:4304-4308.

25

Fein, H.G., Rosen, S.W. and Weintraub, B.D. (1980) Increased glycosylation of serum human chorionic gonadotropin and subunits from eutopic and ectopic sources: comparison with placental and urinary forms. *J Clin Endocrinol Metab* 50, 1111-1120.

30

Fiete, D., V. Srivastava, O. Hindsgaul, and J. U. Baenziger. 1991. *Cell* 67:1103-1110.

35

Grotjan, H. R. J. and L. A. Cole. 1989. In *Microheterogeneity of Glycoprotein Hormones*. H. R. J. Grotjan and L. A. Cole, editors. CRC Press, Boca Raton. 219-237.

Hakim, R. B., R. H. Gray, and H. Zacur. 1995. *Am. J. Obstet. Gynecol.* 172:1510-1517.

5 Hardy, M. R. and R. R. Townsend. 1988. *Proc. Natl. Acad. Sci. U. S. A.* 85:3289-3293.

Hardy, M. R., R. R. Townsend, and Y. C. Lee. 1997. *Anal Biochem* 1988 Apr;170(1):54-62,

10 Hardy, M. R. and R. R. Townsend. 1989. *Carbohydr. Res.* 188:1-7.

15 Hardy, M. R. and R. R. Townsend. 1994. *Methods Enzymol.* 230:208-225.

Harrison, R. F. 1985. *Europ. J. Obstet. Gynec. reprod. Biol.* 20:159-168.

20 Hill, J. A. and D. J. Anderson. 1990. *Archsm. Immun. Ther. Exp.* 38:111-119.

Ho, H-H, O'Connor, JF, Overstreet, JW, Lasley, BL. *Early Pregnancy: Biology and Medicine*, 1997, (in press).

25 Hoermann, R., G. Spoettl, M. Grossmann, B. Saller, and K. Mann. 1997. *J. Clin. Invest.* 71:953-960.

30 Hoermann, R., P. Berger, G. Spoettl, F. Gillesberger, A. Kardana, L. A. Cole, and K. Mann. 1994. *Clin. Chem.* 40:2306-2312.

Hussa, R. O. 1987. *The Clinical Marker hCG*. Praeger, New York.

35 Kagimoto, A., R. Sakakibara, N. Fukushima, N. Ikeda, and M. Ishiguro. 1995. *Biol. Pharm. Bull.* 18:810-817.

001080"ST000000

- Kardana, A., G. D. Braunstein, and L. A. Cole. 1996. *Oncol. Res.* 8:13-16.
- Kardana, A. and L. A. Cole. 1992. *Clin. Chem.* 38:26-33.
- 5
Karlsson, F. A., P. Burman, O. Kampe, J. E. Westlin, and L. Wide. 1993. *Acta Endocrinol. (Copenh)* 129:291-295.
- Kawasaki, T. and G. Ashwell. 1976. *J. Biol. Chem.* 251:1296-1302.
- 10
Kovalevskaya, G, Birken, S, O=Connor, J, Schlatterer, J, Maydelman, Y, Canfield, R. *Endocrine*, 1995, 3:881-887.
- 15
Kumarasamy, R. 1990. *J. Chromatogr.* 512:149-155.
- Lasley, B. L., P. Lohstroh, A. Kuo, E. B. Gold, B. Eskenazi, S. J. Samuels, D. R. Stewart, and J. W. Overstreet. 1995. *Am. J. Ind. Med.* 28:771-781.
- 20
Maack, T., C. H. Park, and M. J. F. Camargo. 1985. In *The kidney: Physiology and pathophysiology*. D. W. Seldin and G. Giebisch, editors. Raven Press, New York. 1773-1803.
- 25
Matzuk, M. M., A. J. Hsueh, P. Lapolt, A. Tsafiriri, J. L. Keene, and I. Boime. 1990. [published erratum appears in *Endocrinology* 1990 Apr;126(4):2204]. *Endocrinology* 126:376-383.
- 30
Moyle, W. R., O. P. Bahl, and L. Marz. 1975. *Journal of Biological Chemistry* 250:9163-9169.
- Newman, C. B., S. L. Wardlaw, and A. G. Frantz. 1985. *Life Sci.* 36:1661-1668.
- 35
Nishimura, R., T. Kitajima, K. Hasegawa, K. Takeuchi,

and M. Mochizuki. 1989. *Jpn. J. Cancer Res.* 80:968-974.

5 Nishimura, R., T. Utsunomiya, K. Ide, T. Kitajima, H. C. Chen, J. L. Strobil, R. O. Husa, and M. Mochizuki. 1987. *Jpn. J. Cancer Res.* 78:833-839.

Nishimura, R., K. Ide, T. Utsunomiya, T. Kitajima, Y. Yuki, and M. Mochizuki. 1988. *Endocrinology* 123:420-425.

10 O'Connor, J., G. Kovalevskaya, S. Birken, J. P. Schlatterer, D. Schechter, D. McMahon, and R. E. Canfield. 1998. *Hum. Reprod.* 13:826-835.

15 O'Connor, J. F., S. Birken, J. W. Lustbader, A. Krichevsky, Y. Chen, and R. E. Canfield. 1994. *Endocr. Rev.* 15:650-683.

20 O'Connor, J. F., J. P. Schlatterer, S. Birken, A. Krichevsky, E. G. Armstrong, D. McMahon, and R. E. Canfield. 1988. *Cancer Res.* 48:1361-1366.

Odell, W. D. and J. Griffin. 1989. *J. Clin. Endocrinol. Metab.* 69:528-532.

25 Odell, W. D. and J. Griffin. 1987. *N. Engl. J. Med.* 317:1688-1691.

Pollak, S., S. Halpine, B. T. Chait, and S. Birken. 1990. *Endocrinology* 126:199-208.

30 Puisieux, A., D. Bellet, F. Troalen, A. Razafindratsita, C. Lhomme, C. Bohuon, and J. M. Bidart. 1990. *Endocrinology* 126:687-694.

35 Quadri, K. H., J. Bernardini, A. Greenberg, S. Laifer, A. Syed, and J. L. Holley. 1994. *Am. J. Kidney Dis.* 24:416-420.

001030"5T20E960

Quenby, S. and R. G. Farquharson. 1994. *Fertil. Steril.* 62:708-710.

5 Ravindranath, N., N. S. Srilatha, M. R. Sairam, and N. R. Moudgal. 1992. *Indian J. Exp. Biol.* 30:982-986.

Rohrer, J. S., G. A. Cooper, and R. R. Townsend. 1993. *Anal. Biochem.* 212:7-16.

10 Sairam, M. R. and L. G. Jiang. 1992. *Mol. Cell Endocrinol.* 85:227-235.

Sairam, M. R. 1989. *FASEB J.* 3:1915-1926.

15 Sairam, M. R., J. Linggen, and G. N. Bhargavi. 1988. *Biosci. Rep.* 8:271-278.

Skarulis, M. C., R. E. Wehmann, B. C. Nisula, and D. L. Blithe. 1992. *J. Clin. Endocrinol. Metab.* 75:91-96.

20 Spellman, M. W. 1990. *Anal. Chem.* 62:1714-1722.

25 Stanton, P. G., G. Poznek, P. G. Burgon, D. M. Robertson, and M. T. W. Hearn. 1993. *J Endocrinol.* 138:529-543.

Szkudlinski, M. W., N. R. Thotakura, I. Bucci, L. R. Joshi, A. Tsai, J. East-Palmer, J. Shiloach, and B. D. Weintraub. 1993. *Endocrinology* 133:1490-1503.

30 Szkudlinski, M. W., N. R. Thotakura, J. E. Tropea, M. Grossmann, and B. D. Weintraub. 1995. *Endocrinology* 136:3325-3330.

35 Taussky, H. H. 1954. *J. Biol. Chem.* 208:853-861.

Thotakura, N. R., M. W. Szkudlinski, and B. D.

007080" ST00E960

Weintraub. 1994. *Glycobiology*. 4:525-533.

Townsend, R. R., M. Hardy, J. D. Olechno, and S. R. Carter. 1988. *Nature* 335:379-380.

5

Townsend, R. R., P. H. Atkinson, and R. B. Trimble. 1991. *Carbohydr. Res.* 215:211-217.

10 Townsend, R. R., M. R. Hardy, and Y. C. Lee. 1989. *Methods Enzymol.* 179:65-76.

Townsend, R. R., M. R. Hardy, D. A. Cumming, J. P. Carver, and B. Bendiak. 1989. *Anal. Biochem.* 182:1-8.

15 Ulloa-Aguirre, A., Mendez, J.P., Cravioto, A., Grotjan, E., Damian-Matsumura, P. and Espinoza, R. (1990) Studies on the microheterogeneity of chorionic gonadotrophin secreted by the human cytotrophoblast in culture. *Hum. Reprod.* 5, 661-669.

20

Weitzhandler, M., D. Kadlecsek, N. Avdalovic, J. G. Forte, D. Chow, and R. R. Townsend. 1993. *J. Biol. Chem.* 268:5121-5130.

25 Wide, L. and B. Hobson. 1987. *Acta Endocrinol. (Copenh)* 116:465-472.

30 Wilcox, A. J., C. R. Weinberg, J. F. O'Connor, D. D. Baird, J. P. Schlatterer, R. E. Canfield, E. G. Armstrong, and B. C. Nisula. 1988. *N. Engl. J. Med.* 319:189-194.

Zinaman, MJ, Clegg, ED, Brown, CC, O'Connor, J, Selevan, SG. *Fertil Steril.*, 1996, 65:503-509.

35

000000 "S" 2000000